



Radial Head/Neck Rehab Non-op Protocol

Dr. Adam V. Metzler

Sports Medicine and Trauma Orthopaedic Surgery

If you have any questions about this protocol or about the patient, do not hesitate to email Dr. Metzler at: ametzler@orthonky.com

Introduction

This protocol is recommended for Mason's type I fracture or type II or III with ORIF.

Goal: Regain full pain-free ROM of elbow and prevent shoulder and wrist stiffness.

Phase (0 to 14 days)

- Begin elbow active ROM and active-assisted ROM for flexion and extension.
- ROM by the end of 2nd week should be at least 15 to 105 degrees.
- Begin putty/grip exercises.
- Begin isometric strengthening exercises for the elbow and wrist.

Phase II (15 days to 6 weeks)

- Continue elbow active and active assisted ROM exercises.
- Full flexion and extension ROM should be achieved by the end of 6 weeks.
- Begin active and active assisted supination and pronation.
- Begin light isotonic strengthening of flexion and extension.
- Assess and maintain shoulder and wrist strength and ROM.

Phase III (7-12 weeks)

- Continue active and active assisted supination and pronation.
- Full pronation and supination should be achieved by the end of the 8th week.
- Progressively increase isotonic strengthening in flexion/extension and pronation/supination.
- Work on any deficits.
- Full return to work around 12-16 weeks based on work level and activity.